

(a) - producing a plurality of amplicates of a section of the [this] nucleic acid with the aid of two primers, one of which can bind to a binding sequence (A) of one strand of the nucleic acid and the other can bind to a binding sequence C' which is essentially complementary to a sequence C which is located in the 3' direction from A and does not overlap A,

(b) - contacting the amplicates with a probe having a binding sequence D which can bind to a sequence B located between the sequences A and C or to the complement thereof, and

(c) - detecting the formation of a hybrid of the amplicate and probe,

wherein the sequence located between the binding sequences A and C contains no nucleotides that do not belong to the sequence region E formed from the binding sequence D of the probe and the sequence of the amplicate bound thereto and the amplicate does not exceed a total length of 100 nucleotides.

2. (Amended) The method of [Method as claimed in] claim 1, wherein the binding sequence D of the probe overlaps one or both binding sequences of the primers.
3. (Amended) The method of claim 1 [Method as claimed in one of the previous claims], wherein at least one of the primers has nucleotides in its non-extendible part which do not hybridize directly with the nucleic acid to be detected or with its complement.
4. (Amended) The method of claim 1 [Method as claimed in one of the previous claims], wherein at least one of the binding sequences is not specific for the nucleic acid to be detected.
5. (Amended) The method of claim 1 [Method as claimed in one of the previous claims], wherein the total length of the amplicates does not exceed 61 nucleotides.

6. (Amended) The method of claim 1 [Method as claimed in one of the previous claims], wherein at least one of the primers is immobilizably-labelled and the probe is detectably-labelled.
7. (Amended) The method of claim 1 [Method as claimed in one of the previous claims], wherein at least one of the primers is detectably-labelled and the probe is immobilizably-labelled or is immobilized.
8. (Amended) The method of claim 1 [Method as claimed in one of the previous claims], wherein the probe is labelled with a fluorescence quencher as well as with a fluorescent dye.
9. (Amended) The method of claim 1 [Method as claimed in one of the previous claims], wherein one of the primers labelled with a first energy transfer component and the probe is labelled with a second energy transfer component which is different from the first energy transfer component.
10. (Amended) The method of claim 1 [Method as claimed in one of the previous claims], wherein the amplificate is detected by physical and/or spectroscopic methods.
11. (Amended) The method of claim 1 [Method as claimed in one of the previous claims], wherein at least one of the primers is not specific for the nucleic acid to be detected.
12. (Amended) The method of [Method as claimed in] claim 11, wherein two of the primers are not specific for the nucleic acid to be detected.
13. (Amended) The method of claim 11 [Method as claimed in one of the claims of 11 and 12], wherein the probe is not specific for the nucleic acid to be detected.

14. (Amended) The method of claim 1 [Method as claimed in one of the previous claims], wherein nucleotides which are each complementary to A, G, C and T are used in the amplification.
15. (Amended) The method of claim 1 [Method as claimed in one of the previous claims], wherein the amplicates are detected by means of mass spectroscopy.
16. (Amended) A method [Method] for the specific detection of a nucleic acid comprising the steps:

(a) - producing a plurality of amplicates of a section of the [this] nucleic acid with the aid of at least two primers,

(b) - contacting the amplicates with a probe which can bind to the amplicate, and

(c) - detecting the formation of a hybrid of the amplicate and the probe,

wherein at least one of the primers is not specific for the group of organisms to which the organism to be detected belongs and the total length of the amplicate does not exceed 100 base pairs.

17. (Amended) The method of [Method as claimed in] claim 16, wherein two of the primers are not specific for the nucleic acid to be detected.
18. (Amended) The method of claim 16 [Method as claimed in one of the claims 16 and 17], wherein the probe is not specific for the nucleic acid to be detected.
19. (Amended) The method of claim 16 [Method as claimed in one of the claims 16 to 18], wherein nucleotides which are each complementary to A, G, C and T are used in the amplification.